

Health and Safety Bulletin #5 December 2004

The Health and Safety/Risk Management Committee would like to make you aware of Head Lice. I'm sure most parents have dealt with this nuisance problem at one time or another. The following is information from www.cdc.gov/ncidod/dpd/parasites/lice/factsheet_head_lice.htm, in response to a minor problem during the 2004 summer season.

It is easy for parents/unit leaders to check their scouts for head lice and we strongly recommend that this be done before/after camping trips and prior to scout camp, if possible. Thank you for helping to insure a safe program for the scouts of the Finger Lakes Council.

What are head lice?


Also called *Pediculus humanus capitis* (peh-DICK-you-lus HUE-man-us CAP-ih-TUS), head lice are parasitic insects found on the heads of people. Having head lice is very common. However, there are no reliable data on how many people get head lice in the United States each year.

Who is at risk for getting head lice?

Anyone who comes in close contact (especially head-to-head contact) with someone who already has head lice is at greatest risk. Occasionally, head lice may be acquired from contact with clothing (such as hats, scarves, coats) or other personal items (such as brushes or towels) that belong to an infested person. Preschool and elementary-age children, 3-11, and their families are infested most often. Girls get head lice more often than boys, women more than men. In the United States, African-Americans rarely get head lice. Personal hygiene or cleanliness in the home or school has nothing to do with getting head lice.

What do head lice look like?

There are three forms of lice: the egg (also called a nit), the nymph, and the adult.



three lice forms compared to a penny (Photo credit:

Egg/Nit: Nits are head lice eggs. They are very small, about the size of a knot in thread, hard to see, and are often confused for dandruff or hair spray droplets. Nits are laid by the adult female at the base of the hair shaft nearest the scalp. They are firmly attached to the hair shaft. They are oval and usually yellow to white. Nits take about 1 week to hatch. Eggs that are likely to hatch are usually located within 1/4 inch of the scalp.




Illustration of egg on a hair shaft
(Image credit: CDC)



Nymph form
(Photo credit: CDC)

Nymph: The nit hatches into a baby louse called a nymph. It looks like an adult head louse, but is smaller. Nymphs mature into adults about 7 days after hatching. To live, the nymph must feed on blood.

Adult: The adult louse is about the size of a sesame seed, has six legs, and is tan to greyish-white. In persons with dark hair, the adult louse will look darker. Females, which are usually larger than the males, lay eggs. Adult lice can live up to 30 days on a person's head. To live, adult lice need to feed on blood. If the louse falls off a person, it dies within 2 days.



Adult louse
(Image credit: CDC)

Where are head lice most commonly found?



Adult louse claws
(Photo credit: CDC)

They are most commonly found on the scalp, behind the ears and near the neckline at the back of the neck. Head lice hold on to hair with hook-like claws found at the end of each of their six legs. Head lice are rarely found on the body, eyelashes, or eyebrows.

What are the signs and symptoms of head lice infestation?

- Tickling feeling of something moving in the hair.
- Itching, caused by an allergic reaction to the bites.
- Irritability.
- Sores on the head caused by scratching. These sores can sometimes become infected.

How did my child get head lice?

Contact with an already infested person is the most common way to get head lice. Head-to-head contact is common during play at school and at home (sports activities, on a playground, slumber parties, at camp).

Less commonly,

- Wearing clothing, such as hats, scarves, coats, sports uniforms, or hair ribbons, recently worn by an infested person.
- Using infested combs, brushes, or towels.
- Lying on a bed, couch, pillow, carpet, or stuffed animal that has recently been in contact with an infested person.

How is head lice infestation diagnosed?

An infestation is diagnosed by looking closely through the hair and scalp for nits, nymphs, or adults. Finding a nymph or adult may be difficult; there are usually few of them and they can move quickly from searching fingers. If crawling lice are not seen, finding nits within a 1/4 inch of the scalp confirms that a person is infested and should be treated. If you only find nits more than 1/4 inch from the scalp (and don't see a nymph or adult louse), the infestation is probably an old one and does not need to be treated. If you are not sure if a person has head lice, the diagnosis should be made by your health care provider, school nurse, or a professional from the local health department or agricultural extension service.



Photo credit: CDC

This fact sheet is for information only and is not meant to be used for self-diagnosis or as a substitute for consultation with a health care provider. If you have any questions about the disease described above or think that you may have a parasitic infection, consult a health care provider.

Revised August 19, 2005